

Gravitational waves during inflation in presence of a decaying cosmological parameter from a 5D vacuum theory of gravity

¹Silvina Paola Gomez Martínez ^{*}, ³José Edgar Madriz Aguilar[†] and ^{1,2}Mauricio Bellini[‡]

¹ *Departamento de Física, Facultad de Ciencias Exactas y Naturales,
Universidad Nacional de Mar del Plata, Punes 3350, (7600) Mar del Plata, Argentina.*

² *Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET).*

³ *Departamento de Física, Universidade Federal da Paraíba,
Caixa Postal 5008, 58059-970 João Pessoa, Pb, Brazil.*

We study gravitational waves generated during the inflationary epoch in presence of a decaying cosmological parameter on a 5D geometrical background which is Riemann flat. Two examples are considered, one with a constant cosmological parameter and the second with a decreasing one.

PACS numbers: 04.20.Jb, 11.10.kk, 98.80.Cq

I. INTRODUCTION

Gravitational waves has been subject of attention since long time ago [1]. Under the standard model of cosmology plus the theory of inflation, it is very natural to predict the existence of the background of gravitational waves [2]. Among the primordial perturbations generated during inflation, there were basically normal scalar part and tensor part. The primordial scalar perturbations provided the seeds of large scale structure which then had gradually formed today's galaxies, which is being tested in current observations of cosmic microwave background (CMB). The tensor perturbations have escaped out of the horizon during inflation, and thus can be completely conserved to form the relic of background gravitational waves, which carries the information of the very early universe. In this sense the tensor perturbations are very significant for studies of very early universe. Their amplitude is related to the energy scale of inflation and they are potentially detectable via observations of B -mode polarization in the cosmic microwave background if the energy scale of inflation is larger than $\sim 3 \times 10^{15}$ GeV [3, 4, 5, 6]. Such a detection would be very important to test inflation. The direct detection of GW is one of the most exciting scientific goals because it would improve our understanding of laws governing the early universe and provide new means to observe it. The most sensitive detectors which are already operating, under construction or being planned, are based on optical interferometers [7]. In particular, gravitational wave perturbations will be measured in future by the Planck satellite, which is designed to produce high-resolution temperature and polarization maps of CMB. If no primordial GW are detected by CMB, a direct-detection experiment to understand the simplest form of inflation must have a sensitivity improved by two to three orders of magnitude over current plans [8]. The basal mechanism of generation of primordial gravitational waves in cosmology has been discussed in [9]. There are two main 4D formalisms developed in the literature; the coordinate based approach of Bardeen [10] and the covariant formalism [11].

The idea that our universe is a 4D space-time embedded in a higher dimensional has been a topic of increased interest in several branches of physics, and in particular, in cosmology. This idea has generated a new kind of cosmological models that includes quintessential expansion. In particular, theories on which is considered only one extra dimension have become quite popular in the scientific community. Among these theories are counted the braneworld scenarios [12], the Space-Time-Matter (STM) theory [13] and all noncompact Kaluza-Klein theories.

In this work we shall study the evolution of gravitational waves on the early universe, which is governed by a (decaying) cosmological parameter $\Lambda(t)$, from a 5D vacuum state defined on a $4 + 1$ Riemann flat spacetime. A decaying cosmological parameter can be introduced in a geometrical manner through the 5D background line element [14]

$$dS_b^2 = \psi^2 \frac{\Lambda(t)}{3} dt^2 - \psi^2 e^2 \int \sqrt{\Lambda/3} dt dr^2 - d\psi^2, \quad (1)$$

where $dr^2 = \delta_{ij} dx^i dx^j$, being $\{x^i\} = \{x, y, z\}$ the local cartesian coordinates. In addition t and ψ are the time-like and fifth space-like local coordinates respectively. Adopting a natural unit system ($\hbar = c = 1$), the fifth coordinate ψ has spatial units whereas the cosmological parameter $\Lambda(t)$ has units of $(length)^{-2}$. The background metric in (1) is

^{*} E-mail address: gomezmar@mdp.edu.ar

[†] E-mail address: jemadriz@fisica.ufpb.br

[‡] E-mail address: mbellini@mdp.edu.ar

Riemann-flat, $R^A{}_{BCD} = 0$, describing perfectly a 5D geometric vacuum.

The usual approach would consist to consider the tensor perturbed line element obtained from (1)

$$dS^2 = \psi^2 \frac{\Lambda(t)}{3} dt^2 - \psi^2 e^{2 \int \sqrt{\Lambda/3} dt} (\delta_{ij} + Q_{ij}) dx^i dx^j - d\psi^2, \quad (2)$$

being $Q_{ij}(t, \vec{r}, \psi)$ the transverse traceless tensor denoting the tensor fluctuations with respect to the metric background (1), and thereby it satisfies $tr(Q_{ij}) = Q^i{}_i = 0$ and $Q_{;i}^{ij} = 0$. The 3D spatial components of the metric (2) can be written as $g_{ij} = -\psi^2 \exp \left[2 \int \sqrt{\Lambda/3} dt \right] (\delta_{ij} + Q_{ij})$, so that its contravariant components can be linearly approximated by $g^{ij} \simeq -\psi^{-2} \exp \left[-2 \int \sqrt{\Lambda/3} dt \right] (\delta^{ij} - Q^{ij})$.

Under this approach the dynamics obeyed by the tensor fluctuations Q_{ij} , is obtained using the 5D linearized Einsteins equations in the 5D vacuum $\delta R_{AB} = 0$. However, as it is well-known solely the space-space components contribute for tensor fluctuations, so the former expression reduces simply to

$$\delta R_{ij} = 0. \quad (3)$$

Hence, after some algebra, one obtain the dynamics of the tensor fluctuations Q_{ij} is determined by

$$\ddot{Q}_{ij} + \left[3\sqrt{\frac{\Lambda}{3}} - \frac{1}{2} \frac{\dot{\Lambda}}{\Lambda} \right] \dot{Q}_{ij} - \frac{\Lambda}{3} e^{-2 \int \sqrt{\Lambda/3} dt} \nabla_r^2 Q_{ij} - \frac{\Lambda}{3} [4\psi Q_{ij,\psi} + \psi^2 Q_{ij,\psi,\psi}] = 0, \quad (4)$$

where $(,)$ denotes the partial derivative while the dot is denoting ordinary derivation with respect the cosmic time t .

II. THE 5D TENSOR MODES

In this letter we shall consider a different approach, which, give us the dynamics (4), but from the the action $I = - \int d^4x d\psi {}^{(5)}L$, given by the background gravitational Lagrangian, plus free tensorial fields, Q_{ij} (A, B run from 0 to 4 and i, j from 1 to 3)

$${}^{(5)}L = \sqrt{\left| \frac{{}^{(5)}g}{{}^{(5)}g_0} \right|} \left[\frac{{}^{(5)}\mathcal{R}}{16\pi G} + \frac{M_p^2}{2} g^{AB} Q_{;A}^{ij} Q_{ij;B} \right], \quad (5)$$

being $(;)$ the covariant derivative. Furthermore, ${}^{(5)}\mathcal{R} = 0$ is the 5D background Ricci scalar and ${}^{(5)}g = \psi^8 \Lambda e^{6 \int \sqrt{\Lambda/3} dt} / 3$ is the determinant of the background covariant metric tensor g_{AB} . Following the usual quantization process for $Q_{ij}(t, \vec{r}, \psi)$ we demand that the next commutation relation must be satisfied

$$\left[Q_{ij}(t, \vec{r}, \psi), \frac{\partial L}{\partial Q_{lm,t}}(t, \vec{r}', \psi') \right] = i \delta_i^l \delta_j^m g^{tt} M_p^2 \sqrt{\left| \frac{{}^{(5)}g}{{}^{(5)}g_0} \right|} \left(\frac{\psi_0}{\psi} \right)^3 e^{-\int [3\sqrt{\frac{\Lambda}{3}} - \frac{\dot{\Lambda}}{2\Lambda}] dt} \times \delta^{(3)}(\vec{r} - \vec{r}') \delta(\psi - \psi'). \quad (6)$$

On the other hand, ${}^{(5)}g_0 \equiv {}^{(5)}g[\psi = \psi_0, \Lambda_0 \equiv \Lambda(t = t_0)]$, being ψ_0 and t_0 some constants to be specified. In this work we are interested to study the dynamics of Q_{ij} . We express the functions Q_{ij} as a Fourier expansion of the form

$$Q^i{}_j(t, \vec{r}, \psi) = \frac{1}{(2\pi)^{3/2}} \int d^3k_r dk_\psi \sum_\alpha {}^{(\alpha)}e^i{}_j \left[a_{k_r k_\psi}^{(\alpha)} e^{i\vec{k}_r \cdot \vec{r}} \zeta_{k_r k_\psi}(t, \psi) + a_{k_r k_\psi}^{(\alpha)\dagger} e^{-i\vec{k}_r \cdot \vec{r}} \zeta_{k_r k_\psi}^*(t, \psi) \right], \quad (7)$$

with α counting the number of polarization degrees of freedom, the asterisk $(*)$ denoting complex conjugate and the creation $a_{k_r k_\psi}^{(\alpha)\dagger}$ and annihilation $a_{k_r k_\psi}^{(\alpha)}$ operators satisfying the algebra

$$\left[a_{k_r k_\psi}^{(\alpha)}, a_{k'_r k'_\psi}^{(\alpha')\dagger} \right] = g^{\alpha\alpha'} \delta^{(3)}(\vec{k}_r - \vec{k}'_r) \delta(\psi - \psi'), \quad \left[a_{k_r k_\psi}^{(\alpha)}, a_{k'_r k'_\psi}^{(\alpha')} \right] = \left[a_{k_r k_\psi}^{(\alpha)\dagger}, a_{k'_r k'_\psi}^{(\alpha')\dagger} \right] = 0. \quad (8)$$

The polarization tensor ${}^{(\alpha)}e_{ij}$ obeys

$${}^{(\alpha)}e_{ij} = {}^{(\alpha)}e_{ji}, \quad {}^{(\alpha)}e_{ii} = 0, \quad k^i {}^{(\alpha)}e_{ij} = 0, \quad {}^{(\alpha)}e_{ij}(-\vec{k}_r) = {}^{(\alpha)}e_{ij}^*(\vec{k}_r). \quad (9)$$

Inserting (7) in (4) we obtain that the dynamics of the 5D tensor modes $\zeta_{k_r m}(t, \psi)$ is given by

$$\ddot{\zeta}_{k_r k_\psi} + \left[3\sqrt{\frac{\Lambda}{3}} - \frac{1}{2} \frac{\dot{\Lambda}}{\Lambda} \right] \dot{\zeta}_{k_r k_\psi} + \left[\frac{\Lambda}{3} k_r^2 e^{-2 \int \sqrt{\Lambda/3} dt} - \frac{\Lambda}{3} \left(4\psi \frac{\partial}{\partial \psi} + \psi^2 \frac{\partial^2}{\partial \psi^2} \right) \right] \zeta_{k_r k_\psi} = 0. \quad (10)$$

Decomposing the tensor modes $\zeta_{k_r k_\psi}(t, \psi)$ into Kaluza-Klein modes

$$\zeta_{k_r k_\psi}(t, \psi) \sim \xi_{k_r}(t) \Theta_m(\psi), \quad (11)$$

equation (10) yields

$$\ddot{\xi}_{k_r} + \left(3\sqrt{\frac{\Lambda}{3}} - \frac{1}{2} \frac{\dot{\Lambda}}{\Lambda} \right) \dot{\xi}_{k_r} + \left[\frac{\Lambda}{3} e^{-2 \int \sqrt{\Lambda/3} dt} k_r^2 + m^2 \frac{\Lambda}{3} \right] \xi_{k_r} = 0, \quad (12)$$

$$\psi^2 \frac{d^2 \Theta_m}{d\psi^2} + 4\psi \frac{d\Theta_m}{d\psi} + m^2 \Theta_m = 0, \quad (13)$$

where the parameter $m^2 = (k_\psi \psi)^2$ is related with the squared of the KK-mass measured by a class of observers in 5D. Using the transformation $\xi_{k_r}(t) = \exp[-(1/2) \int (3\sqrt{\Lambda/3} - (\dot{\Lambda}/2\Lambda)) dt] \chi_{k_r}(t)$ and $\Theta_m(z) = e^{-3/2 z} L_m(z)$, with $z = \ln(\psi/\psi_0)$, in eqs. (12) and (13), respectively, we obtain

$$\ddot{\chi}_{k_r} + \left[\frac{\Lambda}{3} e^{-2 \int \sqrt{\Lambda/3} dt} k_r^2 - \frac{1}{4} \sqrt{\frac{3}{\Lambda}} \dot{\Lambda} + \frac{1}{4} \frac{\ddot{\Lambda}}{\Lambda} - \frac{5}{16} \frac{\dot{\Lambda}^2}{\Lambda^2} + \frac{3}{4} \sqrt{\frac{\Lambda}{3}} \frac{\dot{\Lambda}}{\Lambda} + \left(\frac{m^2}{3} - \frac{3}{4} \right) \Lambda \right] \chi_{k_r} = 0, \quad (14)$$

$$\frac{d^2 L_m(z)}{dz^2} + \left[m^2 - \frac{9}{4} \right] L_m(z) = 0. \quad (15)$$

This way, given a cosmological parameter $\Lambda(t)$, the temporal evolution of the tensor modes $\xi_{k_r}(t)$ in 5D is determined by solutions of (14). Once a solution for $\xi_{k_r}(t)$ is obtained, it should satisfy the algebra (6). This can be made guaranteeing that such a solution obey

$$\chi_{k_r} \dot{\chi}_{k_r}^* - \dot{\chi}_{k_r} \chi_{k_r}^* = i, \quad |L_m|^2 = 1. \quad (16)$$

On the other hand, note that equation (13) is exactly the same as the one obtained in [15]. Therefore about the behavior of the modes with respect the fifth coordinate we can say that for $m > 3/2$ the KK-modes are coherent on the ultraviolet sector (UV), described by the modes

$$k_r^2 > \left\{ \frac{3}{2\Lambda} \frac{d}{dt} \left[3\sqrt{\frac{\Lambda}{3}} - \frac{\dot{\Lambda}}{2\Lambda} \right] - \frac{3}{4\Lambda} \left(3\sqrt{\frac{\Lambda}{3}} - \frac{\dot{\Lambda}}{2\Lambda} \right)^2 - m^2 \right\} e^{2 \int \sqrt{\Lambda/3} dt} > 0. \quad (17)$$

However, for $m < 3/2$ those modes are unstable and diverge at infinity. The modes with $m = 0$ and $m > 3/2$ comply with the conditions (16), so that they are normalizable.

III. EFFECTIVE 4D DYNAMICS

To describe the 4D dynamics we can make a foliation on $\psi = \psi_0$ on the line element (1), such that the effective 4D background metric holds: $dS^2|_{eff} = ds^2$, where

$$ds^2 = \psi_0^2 \frac{\Lambda(t)}{3} dt^2 - \psi_0^2 e^{2 \int \sqrt{\Lambda/3} dt} dr^2. \quad (18)$$

In this section we shall study the dynamics of the 4D tensor-fluctuations $h_{ij}(t, \vec{r}) \equiv Q_{ij}(t, \vec{r}, \psi = \psi_0)$, making emphasis on the long wavelength section, which describes this field on cosmological scales. The effective 4D action $^{(4)}I$ is (α, β run from 0 to 3)

$$^{(4)}I = - \int d^4x \sqrt{\left| \frac{^{(4)}g}{^{(4)}g_0} \right|} \left[\frac{^{(4)}\mathcal{R}}{16\pi G} + \frac{M_p^2}{2} g^{\alpha\beta} Q_{;\alpha}^{ij} Q_{ij;\beta} \right] \Big|_{\psi=\psi_0}, \quad (19)$$

where ${}^{(4)}\mathcal{R} = 12/\psi_0^2$ is the effective 4D Ricci scalar valued on the metric (18), such that we obtain an equation of state which describes an effective 4D vacuum: $p = -\rho$, being p and ρ the pressure and the energy density, respectively. Hence, the metric (1) could be considered as an extension of the Ponce de Leon one[16], which, on a foliation $\psi = \psi_0$, also describes an effective 4D vacuum dominated expansion.

The effective 4D equation of motion for the 4D tensor-fluctuations is

$$\ddot{h}_j^i + \left[3\sqrt{\frac{\Lambda}{3}} - \frac{\dot{\Lambda}}{2\Lambda} \right] \dot{h}_j^i - \frac{\Lambda}{3} e^{-2\int \sqrt{\frac{\Lambda}{3}} dt} \nabla_r^2 h_j^i - \frac{\Lambda}{3} \left[\frac{4}{\psi} \frac{\partial Q_j^i}{\partial \psi} + \psi^2 \frac{\partial^2 Q_j^i}{\partial \psi^2} \right] \Big|_{\psi=\psi_0} = 0. \quad (20)$$

Using the eq. (13), we obtain

$$\ddot{h}_j^i + \left[3\sqrt{\frac{\Lambda}{3}} - \frac{\dot{\Lambda}}{2\Lambda} \right] \dot{h}_j^i - \frac{\Lambda}{3} e^{-2\int \sqrt{\frac{\Lambda}{3}} dt} \nabla_r^2 h_j^i + \frac{m^2 \Lambda}{3} h_j^i = 0. \quad (21)$$

After make the transformation $h_j^i(t, \vec{r}) = e^{-1/2 \int [3(\frac{\Lambda}{3})^{1/2} - \frac{\dot{\Lambda}}{2\Lambda}] dt} \chi_j^i(t, \vec{r})$, we obtain

$$\ddot{\chi}_j^i - \frac{\Lambda}{3} e^{-2\int \sqrt{\frac{\Lambda}{3}} dt} \nabla_r^2 \chi_j^i - \left[\frac{1}{4} \sqrt{\frac{3}{\Lambda}} \dot{\Lambda} + \frac{1}{4} \frac{\ddot{\Lambda}}{\Lambda} - \frac{5}{16} \frac{\dot{\Lambda}^2}{\Lambda^2} + \frac{3}{4} \sqrt{\frac{\Lambda}{3}} \frac{\dot{\Lambda}}{\Lambda} + \left(\frac{m^2}{3} - \frac{3}{4} \right) \Lambda \right] \chi_j^i = 0, \quad (22)$$

such that it is possible to make a Fourier expansion for χ_j^i

$$\chi_j^i(t, \vec{r}) = \frac{1}{(2\pi)^{3/2}} \int d^3 k_r dk_\psi \sum_{\alpha=+, \times} {}^{(\alpha)} e^{i j} \left[a_{k_r k_\psi}^{(\alpha)} e^{i \vec{k}_r \cdot \vec{r}} \chi_{k_r k_\psi}(t, \psi) + a_{k_r k_\psi}^{(\alpha) \dagger} e^{-i \vec{k}_r \cdot \vec{r}} \chi_{k_r k_\psi}^*(t, \psi) \right] \delta(k_\psi - k_{\psi_0}), \quad (23)$$

where we require that the modes $\chi_{k_r} \equiv \chi_{k_r k_{\psi_0}}$ satisfy the commutation relation

$$[\chi_{k_r}(t, \vec{r}), \dot{\chi}_{k_r}(t, \vec{r}')] = i \delta^{(3)}(\vec{r} - \vec{r}'). \quad (24)$$

Using (23) this expression reads

$$\chi_{k_r} \dot{\chi}_{k_r}^* - \chi_{k_r}^* \dot{\chi}_{k_r} = i, \quad (25)$$

which is the condition for the modes to be normalizable on the UV-sector. Inserting (23) in (22) we obtain the dynamical equation for the k_r -modes

$$\ddot{\chi}_{k_r} + \left[\frac{\Lambda}{3} e^{-2\int \sqrt{\frac{\Lambda}{3}} dt} k_r^2 - \left(\frac{1}{4} \sqrt{\frac{3}{\Lambda}} \dot{\Lambda} + \frac{1}{4} \frac{\ddot{\Lambda}}{\Lambda} - \frac{5}{16} \frac{\dot{\Lambda}^2}{\Lambda^2} + \frac{3}{4} \sqrt{\frac{\Lambda}{3}} \frac{\dot{\Lambda}}{\Lambda} + \left(\frac{m^2}{3} - \frac{3}{4} \right) \Lambda \right) \right] \chi_{k_r} = 0. \quad (26)$$

This way for a given $\Lambda(t)$ corresponds a normalized solution for the k_r -modes by solving (26). Once obtained a normalized solution of (26), we will be able of dealing with the spectrum on super Hubble scales. The amplitude of the 4D tensor metric fluctuations $\langle h^2 \rangle = \langle 0 | h^i_j h_i^j | 0 \rangle$ on the IR-sector is given by

$$\langle h^2 \rangle = \frac{4}{\pi^2} e^{-\int [3(\frac{\Lambda}{3})^{1/2} - \frac{\dot{\Lambda}}{2\Lambda}] dt} \int_0^{\epsilon k_H} \frac{dk_r}{k_r} k_r^3 [\chi_{k_r}(t) \chi_{k_r}^*(t)]_{IR}, \quad (27)$$

where $\epsilon = k_{max}^{IR}/k_p \ll 1$ is a dimensionless parameter, being $k_{max}^{IR} = k_H(t_i)$ the wave-number related to the Hubble radius at the time t_i . This time corresponds at the time when the gravitational modes re-enter to the horizon. In addition, k_p is the Planckian wave-number. Clearly, in order to obtain an explicit spectrum it is necessary to specify first a functional form for $\Lambda(t)$. Some illustrative examples will be studied in the next section.

IV. EXAMPLES

In order to illustrate the formalism developed in the previous section, along the present section we study a pair of interesting examples. The first one contemplating a constant cosmological parameter $\Lambda = 3H_0^2$, and the second one considering a decaying $\Lambda(t) = 3p^2/t^2$.

A. Case $\Lambda = 3H_0^2$

The simplest example results of taking the cosmological parameter Λ to be a constant, and in particular $\Lambda = 3H_0^2$. In this particular case the equation of motion for the modes χ_{k_r} becomes

$$\ddot{\chi}_{k_r} + \left[H_0^2 e^{-2H_0 t} k_r^2 - \left(m^2 - \frac{9}{4} \right) H_0^2 \right] \chi_{k_r} = 0. \quad (28)$$

The general solution for this equation is

$$\chi_{k_r}(t) = A_1 \mathcal{H}_\nu^{(1)} [k_r e^{-H_0 t}] + A_2 \mathcal{H}_\nu^{(2)} [k_r e^{-H_0 t}], \quad (29)$$

where A_1 and A_2 are integration constants. After the Bunch-Davies normalization [17] the normalized solution reads

$$\chi_{k_r}(t) = \frac{i}{2} \sqrt{\frac{\pi}{H_0}} \mathcal{H}_\nu^{(2)} [k_r e^{-H_0 t}], \quad (30)$$

with $\nu = (1/2)\sqrt{4m^2 - 9}$. This solution is stable for $m^2 > 9/4$, on scales $k_r^2 > (m^2 - \frac{9}{4}) e^{2H_0 t} > 0$. Now considering the asymptotic expansion for the Hankel function $\mathcal{H}_\nu^{(2)}[y] \simeq (-i/\pi)\Gamma(\nu)[y/2]^{-\nu}$ in (30), the amplitude of the 4D tensor metric fluctuations (27) on cosmological (super Hubble) scales gives

$$\langle h^2 \rangle_{SH} = \frac{2^{2\nu}}{\pi^3(3-2\nu)} \frac{\Gamma^2(\nu)}{H_0} e^{-(3-2\nu)H_0 t} \epsilon^{3-2\nu} k_H^{3-2\nu}, \quad (31)$$

where $k_H(t) \sim e^{H_0 t}$. Hence, the gravitational spectrum $\mathcal{P}_g(k_r)$ takes the form

$$\mathcal{P}_g(k_r) = \frac{2^{2\nu}}{\pi^3} \frac{\Gamma^2(\nu)}{H_0} e^{-(3-2\nu)H_0 t} k_r^{3-2\nu} \Big|_{k_r = \epsilon k_H} \quad (32)$$

We can see from (32) that for $m \simeq 3/\sqrt{2} > (3/2)$, the gravitational spectrum $\mathcal{P}_\nu(k_r)$ is nearly scale invariant and consequently the tensor spectral index becomes $n_T \equiv 3 - 2\nu \simeq 0$ in this case.

B. Case $\Lambda = 3p^2/t^2$

Another interesting case appears considering a decaying cosmological parameter $\Lambda = 3p^2/t^2$, with the restriction $\dot{\Lambda} < 0$. In this case the equation of motion for the modes $\chi_{k_r}(t)$ results to be

$$\ddot{\chi}_{k_r} + \left\{ k_r^2 p^2 t_0^{2p} t^{-2(p+1)} - \left[\left(m^2 - \frac{9}{4} \right) p^2 - \frac{9}{4} p + \frac{1}{4} \right] t^{-2} \right\} \chi_{k_r} = 0, \quad (33)$$

where $M^2(t) = \left[\left(m^2 - \frac{9}{4} \right) p^2 - \frac{9}{4} p + \frac{1}{4} \right] t^{-2}$ can be interpreted as an effective squared term of mass. The permitted values for m should be

$$\frac{9}{4} < m^2 \leq \frac{9}{4} \left(\frac{9}{4} + 1 \right), \quad (34)$$

for which

$$\frac{9 - \sqrt{117 - 16m^2}}{2(4m^2 - 9)} \leq p \leq \frac{9 + \sqrt{117 - 16m^2}}{2(4m^2 - 9)}. \quad (35)$$

The general solution of (33) can be expressed in terms of Bessel functions as

$$\chi_{k_r}(t) = B_1 \left[\frac{y(t)}{2} \right]^{-\mu/(2p)} t^{(1-\mu)/2} \Gamma \left(1 + \frac{\mu}{2p} \right) \mathcal{J}_\mu[y(t)] + B_2 \left[\frac{y(t)}{2} \right]^{\mu/(2p)} t^{(1+\mu)/2} \Gamma \left(1 - \frac{\mu}{2p} \right) \mathcal{J}_{-\mu}[y(t)], \quad (36)$$

where $\mu = \sqrt{4p^2 m^2 - 9p(p+1) + 2}$ and $y(t) = k_r(t_0/t)^p$. In general this expression is not normalizable. However there exist some particular solutions of (35) that are normalizable. A particular case that yields a normalizable solution

results by taking $m = \pm[1/(2p)]\sqrt{9p(p+1)-1}$. In this case the dynamical equation for the quantum gravitational modes reduces to

$$\ddot{\chi}_{k_r} + k_r^2 p^2 t_0^{2p} t^{-2(p+1)} \chi_{k_r} = 0, \quad (37)$$

whose general solution is

$$\chi_{k_r}(t) = \sqrt{t} \left\{ C_1 \mathcal{H}_\omega^{(1)} \left[k_r \left(\frac{t_0}{t} \right)^p \right] + C_2 \mathcal{H}_\omega^{(2)} \left[k_r \left(\frac{t_0}{t} \right)^p \right] \right\}, \quad (38)$$

being C_1 and C_2 integration constants, and $\omega = 1/(2p)$. The Bunch-Davies normalized solution is then

$$\chi_{k_r}(t) = \frac{i}{2} \sqrt{\frac{1}{\pi p}} \mathcal{H}_\omega^{(2)} \left[k_r \left(\frac{t_0}{t} \right)^p \right]. \quad (39)$$

In this case the amplitude of the 4D tensor metric fluctuations on super-Hubble scales ($k_r \gg k_H$) reads

$$\langle h^2 \rangle_{SH} = \frac{2^{2\omega}}{p \pi^5} \frac{\Gamma^2(\omega)}{3-2\omega} \left(\frac{t_0}{t} \right)^{(3-2\omega)p+1} \epsilon^{3-2\omega} k_H^{3-2\omega}. \quad (40)$$

Therefore the gravitational spectrum $\mathcal{P}_g(k_r)$ is in this case

$$\mathcal{P}_g(k_r) = \frac{2^{2\omega}}{p \pi^5} \Gamma^2(\omega) \left(\frac{t_0}{t} \right)^{(3-2\omega)p+1} k_r^{3-2\omega} \Big|_{k_r=\epsilon k_H}. \quad (41)$$

Clearly, for $p \simeq 1/3$ (that corresponds to $m \simeq 3\sqrt{3}/2 > 3/2$), the spectrum is nearly scale invariant i.e. $n_T \equiv 3-2\omega \simeq 0$.

V. FINAL COMMENTS

In this letter we have studied the emergence of gravitational waves in the early universe, which is considered as dominated by a decaying cosmological parameter, from a 5D Riemann flat background metric, on which we define the 5D vacuum. Our approach is different to others, because we consider gravitational waves as originated by a physical field Q_{ij} , but not a tensorial linearized fluctuation of the metric. Therefore, it is possible to deal with Q_{ij} as a quantum traceless tensorial field with null divergence.

The effective 4D dynamics of GW, $h_{ij} = Q_{ij}(\psi = \psi_0)$, can be viewed as induced by the foliation of the fifth coordinate. In this letter we have worked two examples: **a)** In the case with constant cosmological parameter, $\Lambda = \Lambda_0$, we obtain that the KK mass of the gravitons should be $m \simeq 3/\sqrt{2}$ to obtain a nearly scale invariant tensorial power spectrum of $\langle h^2 \rangle_{SH}$. This value of m corresponds to an effective 4D mass $M \simeq 3H_0/2 = \sqrt{3\Lambda}/2$. Note that this mass can be related to the Einstein mass m_E , as it is considered in[18]: $M \simeq \frac{3}{2Gm_E}$. **b)** The case where the cosmological parameter decrease with time is the more interesting. In this case the gravitons are normalizable only when their effective 4D mass M become null. We obtain that the spectrum of the squared fluctuations of gravitational waves are scale invariant for KK masses with values $m \simeq 3\sqrt{3}/2$.

Acknowledgements

SPGM acknowledges UNMdP for financial support. MB acknowledges CONICET and UNMdP (Argentina) for financial support. JEMA acknowledges CNPq-CLAF and UFPB for financial support.

-
- [1] R. Arnowitt and S. Deser, Phys. Rev. **113**, 745 (1959); R. Arnowitt, S. Deser and c. W. Misner, Phys. Rev. **121**, 1556 (1961); T. A. Barnebey, Phys. Rev. **D10**, 1741 (1974); D. L. Lee, Phys. Rev. **D10**, 2374 (1974).
 - [2] A. Starobinsky, JETP Lett. **30**, 682 (1979); V. Rubakov, M. Sazhin and A. Veryaskin, Phys. Lett. **BB115**, 189 (1982); R. Fabbri and M. Pollock, Phys. Lett. **B125**, 445 (1983); L. Abbott and D. Harari, Nucl. Phys. **B264**, 487 (1986); E. Stewart and D. Lyth, Phys. Lett. **B302**, 171 (1983).
 - [3] M. Kamionkowski, A. Kosowski and A. Stebbins, Phys. Rev. **D55**, 7368 (1997); *ibid.*, Phys. Rev. Lett. **78**, 2058 (1997).

- [4] M. Kesden, A. Cooray and M. Kamionkowski, Phys. Rev. Lett. **89**, 011304 (2002).
- [5] L. Knox and Y. Song, Phys. Rev. Lett. **89**, 011303 (2002).
- [6] U. Seljak and C. M. Hirata, Astrophys. J. **463**, 1 (1996).
- [7] For a general update see the following special issue: Class. Quant. Grav. **23** (2006).
- [8] T. L. Smith, H. V. Peiris and A. Cooray, Phys. Rev. **D73**, 123503 (2006).
- [9] L. P. Grinshchuk. Sov. Phys. JETP **40**, 409 (1975); B. Allen, Phys. Rev. **D37**, 2078 (1988).
- [10] J. Bardeen, Phys. Rev. **D22**, 1882 (1980).
- [11] G. F. R. Ellis and M. Bruni, Phys. Rev. **D40**, 1804 (1989).
- [12] L. Randall and R. Sundrum, Phys. Rev. Lett. **83**, 4690 (1999).
- [13] P. S. Wesson, Gen. Rel. Grav. **16**, 193 (1984); P. Wesson, Gen. Rel. Grav. **22**, 707 (1990); P. S. Wesson, Phys. Lett. **B276**, 299 (1992); P. S. Wesson and J. Ponce de Leon, J. Math. Phys. **33**, 3883 (1992); H. Liu and P. S. Wesson, J. Math. Phys. **33**, 3888 (1992); P. Wesson, H. Liu and P. Lim, Phys. Lett. **B298**, 69 (1993).
- [14] M. Bellini, Phys. Lett. **B632** (2006) 610-616.
- [15] J.E. Madriz Aguilar, Phys. Lett. **B645** (2007) 6-11.
- [16] J. Ponce de Leon, Gen. Rel. Grav. **20**, 539 (1988).
- [17] T. S. Bunch and P. C. W. Davies, Proc. Roy. Soc. **A360**, 117 (1978).
- [18] P. S. Wesson, Mod. Phys. Lett. **A19**, 1995 (2004).